

PCTWORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁵ : A61K 31/57, 31/565, 31/58, 9/00, 9/70, C07J 43/00, 1/00, 17/00		A2	(11) International Publication Number: WO 94/16709 (43) International Publication Date: 4 August 1994 (04.08.94)
(21) International Application Number: PCT/CA94/00022 (22) International Filing Date: 19 January 1994 (19.01.94) (30) Priority Data: 08/005,619 19 January 1993 (19.01.93) US 08/180,361 18 January 1994 (18.01.94) US (71) Applicant: ENDORECHERCHE INC. [CA/CA]; 2989 de la Promenade Ste-Foy, Quebec City, Quebec G1W 2J6 (CA). (72) Inventor: LABRIE, Fernand; 2989 de la Promenade Ste-Foy, Quebec City, Quebec G1W 2J6 (CA). (74) Agent: MITCHELL, Richard, J.; Marks & Clerk, P.O. Box 957, Station B, Ottawa, Ontario K1P 5S7 (CA).		(81) Designated States: AT, AU, BB, BG, BR, BY, CA, CH, CN, CZ, DE, DK, ES, FI, GB, HU, JP, KP, KR, KZ, LK, LU, MG, MN, MW, NL, NO, NZ, PL, PT, RO, RU, SD, SE, SK, UA, VN, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG). Published <i>Without international search report and to be republished upon receipt of that report.</i>	
(54) Title: THERAPEUTIC USES AND DELIVERY SYSTEMS OF DEHYDROEPIANDROSTERONE (57) Abstract <p>Sex steroid precursors such as dehydroepiandrosterone and dehydroepiandrosterone sulphate, and compounds converted <i>in vivo</i> to either of the foregoing, are utilized for the treatment and/or prevention of vaginal atrophy, hypogonadism, diminished libido, osteoporosis, urinary incontinence, ovarian cancer, uterine cancer, skin atrophy, for contraception, and, in combination with an estrogen and/or progestin, for the treatment of menopause. The precursors may be formulated for percutaneous or transmucosal administration. Gels, solutions, lotions, creams, ointments and transdermal patches for the administration of these presursors are provided, as are certain pharmaceutical compositions and kits which can be used for the prevention and treatment of a wide variety of conditions related to decreased secretion of sex steroid precursors by the adrenals.</p>			

BEST AVAILABLE COPY

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	GB	United Kingdom	MR	Mauritania
AU	Australia	GE	Georgia	MW	Malawi
BB	Barbados	GN	Guinea	NE	Niger
BE	Belgium	GR	Greece	NL	Netherlands
BF	Burkina Faso	HU	Hungary	NO	Norway
BG	Bulgaria	IE	Ireland	NZ	New Zealand
BJ	Benin	IT	Italy	PL	Poland
BR	Brazil	JP	Japan	PT	Portugal
BY	Belarus	KE	Kenya	RO	Romania
CA	Canada	KG	Kyrgyzstan	RU	Russian Federation
CF	Central African Republic	KP	Democratic People's Republic of Korea	SD	Sudan
CG	Congo	KR	Republic of Korea	SE	Sweden
CH	Switzerland	KZ	Kazakhstan	SI	Slovenia
CI	Côte d'Ivoire	LJ	Liechtenstein	SK	Slovakia
CM	Cameroon	LK	Sri Lanka	SN	Senegal
CN	China	LU	Luxembourg	TD	Chad
CS	Czechoslovakia	LV	Latvia	TG	Togo
CZ	Czech Republic	MC	Monaco	TJ	Tajikistan
DE	Germany	MD	Republic of Moldova	TT	Trinidad and Tobago
DK	Denmark	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	US	United States of America
FI	Finland	MN	Mongolia	UZ	Uzbekistan
FR	France			VN	Viet Nam
GA	Gabon				

THERAPEUTIC USES AND DELIVERY SYSTEMS OF DEHYDROEPIANDROSTERONE

5

10

BACKGROUND OF THE INVENTION

15 This invention relates to a method for preventing and/or treating vaginal atrophy, hypogonadism, diminished libido, osteoporosis, urinary incontinence, ovarian cancer, uterine cancer, and menopause or contraception in susceptible warm-blooded animals including humans involving administration of dehydroepiandrosterone (DHEA), dehydroepiandrosterone-sulfate (DHEA-S) or compounds converted *in vivo* to either and to pharmaceutical products, 20 including kits and pharmaceutical compositions for delivery of active ingredient(s) useful to the invention.

Primates are unique in having adrenals that secrete large amounts of the precursor steroid dehydroepiandrosterone (DHEA) and especially DHEA-sulfate (DHEA-S), which are converted into androstenedione (D4-dione) or 25 androstene-diol (D⁵-diol) and then into potent androgens and estrogens in peripheral tissues (Adams, Mol. Cell. Endocrinol. 41: 1-17, 1985; Labrie et al., in Important Advances in Oncology (de Vita S, Hellman S, Rosenberg SA, eds), JB Lippincott, Philadelphia PA, pp 193-200, 1985). DHEA-S, the major steroid present in blood of both men and women is converted into DHEA and 30 D5-diol in peripheral tissues, thus maintaining a close correlation between the concentration of these three steroids in the blood (Adams, Mol. Cell. Endocrinol. 41: 1-17, 1985). Depending upon the relative activities of 17b-hydroxysteroid dehydrogenase (17b-HSD), aromatase and 5a-reductase, 35 DHEA or its derivatives will be preferentially converted into androgens and/or estrogens.

SUBSTITUTE SHEET

- 2 -

The low serum values of DHEA and DHEA-S found at birth persist up to six years of age. Usually, during the 7th year of age, serum levels of these two steroids increase and continue to rise until age 16 in both boys and girls (Orentreich et al., J. Clin. Endocr. Metab. 59: 551-555, 1984). A further increase is then seen in males, who typically reach maximal levels between 20 and 24 years of age. In women, there is usually no further increase after 16 years. DHEA and DHEA-S decrease with aging in both men and women (Vermeulen and Verdoreck, J. Steroid Biochem. 7: 1-10, 1976; Vermeulen et al., J. Clin. Endocr. Metab. 54: 187-191, 1982). In fact, at 70 years of age, serum DHEA-S levels are at approximately 20% of their peak values while they decrease by up to 95% by the age of 85 to 90 years (Migeon et al., J. Clin. Endocr. Metab. 17: 1051-1062, 1957). The 70% to 95% reduction in the formation of DHEA-S by the adrenals during aging results in a dramatic reduction in the formation of androgens and estrogens in peripheral target tissues, thus resulting in a marked decrease in the biochemical and cellular functions induced by sex steroids.

In addition to the decrease in DHEA-S formation, in men, a progressive decrease in the concentration of testosterone in the spermatic vein (Giusti et al., Exp. Gerontol. 10: 241-245, 1975) as well as in plasma (Lewis et al., Acta Endocrinol. 82: 444-448, 1976; Zumoff et al., J. Clin. Endocr. Metab. 54: 534-538, 1982) has been observed after the age of 60-70 years. Such data, however, have been the subject of controversy (Nieschlag et al., J. Clin. Endocr. Metab. 55: 676-681, 1982). In postmenopausal women, serum testosterone levels are lower than during reproductive life (Forest MG, Physiological changes in circulating androgens, in Androgens in Childhood (Forest MG, ed), Karger Basel, p. 104-129, 1989).

The skin is an important site of sex steroid formation and its function is known to be regulated by sex steroids. Sex steroids can act directly in the skin or may stimulate growth hormone and prolactin secretion by the anterior pituitary gland. Skin atrophy is in fact known to occur in growth hormone deficiency, probably through a secondary decrease in insulin-like growth factor (IGF-1) secretion. Serum growth hormone (GH) and insulin-like growth factor (IGF-1) levels are known to be reduced in aging men and women.

SUBSTITUTE SHEET

- 3 -

Plasma DHEA-S concentration has been suggested as a predictor of osteoporosis (Nordin et al., J. Clin. Endocr. Metab. 60: 651-657, 1985; Deutsch et al., Int. J. Gynecol. Obstet. 25: 217-220, 1987). Serum DHEA has
5 in fact been found to be significantly lower in osteoporotic compared to normal subjects (Nordin et al., J. Clin. Endocr. Metab. 60: 651-657, 1985). Low tissue $\Delta 4$ -dione is also likely to accompany low DHEA-S levels. Since $\Delta 4$ -dione is the precursor of estrone which is a main source of estradiol in postmenopausal women (Marshall et al., Clin. Endocrinol. 9: 407, 1978), such
10 secondary low levels of estrogens are likely to be involved in osteoporosis (Nordin et al., Lancet 2: 277, 1981).

As another mechanism, low serum DHEA-S levels resulting in low formation of androgens in peripheral tissues could also result in low bone formation, a
15 characteristic of postmenopausal osteoporosis (Meunier et al., in Histological heterogeneity of apparently idiopathic osteoporosis and treatment (DeLuca HF, Frost HM, Jee WSS, Johnston Jr CC, Parfitt AM, eds), University Park Press, Baltimore, p. 293). In fact, Deutsch et al. (Int. J. Gynecol. Obstet. 25: 217-222, 1987) have found a significant correlation between serum DHEA-S
20 and androstenedione levels and osteoporosis in late postmenopausal women while no correlation was found between serum estrogens and bone density, thus suggesting a major importance of androgens in postmenopausal bone loss.

25 Studies in animals have shown that androgen deficiency leads to osteopenia while testosterone administration increases the overall quantity of bone (Silverberg and Silverberg, 1971; See Finkelstein et al., Ann. Int. Med. 106: 354-361, 1987). Orchiectomy in rats can cause osteoporosis detectable within 2 months (Winks and Felts, Calcif. Tissue. Res. 32: 77-82, 1980;
30 Verhas et al., Calcif. Tissue Res. 39: 74-77, 1986).

As mentioned earlier, adrenal androgen levels have been found to be reduced in osteoporosis (Nordin et al., J. Clin. Endocr. Metab. 60: 651, 1985). Moreover, elevated androgens in postmenopausal women have been shown
35 to protect against accelerated bone loss (Deutsch et al., Int. J. Gynecol. Obstet. 25: 217-222, 1987; Aloia et al., Arch. Int. Med. 143: 1700-1704,

SUBSTITUTE SHEET

1983). In agreement with such a role of androgens, urinary levels of androgen metabolites are lower in postmenopausal symptomatic menopause than in matched controls and a significant decrease in conjugated dehydroepiandrosterone (DHEA) was found in the plasma of osteoporotic patients (Hollo and Feher, Acta Med. Hung. 20: 133, 1964; Urist and Vincent, J. Clin. Orthop. 18: 199, 1961; Hollo et al., Acta Med. Hung. 27: 155, 1970). It has been suggested that postmenopausal osteoporosis results from both hypoestrogenism and hypoandrogenism (Hollo et al., Lancet, 1357, 1976). Since aging (and osteoporosis) is accompanied by a decrease in an almost indefinite number of parameters and each tissue responds differently (including no response) depending upon the steroidogenic enzymes present, no correlation could be made between changes in DHEA levels and bone loss prior to the present applicants' findings described below regarding bone cells' processing of DHEA.

As a mechanism for the above-suggested role of both estrogens and androgens in osteoporosis, the presence of estrogen (Komm et al., Science 241: 81-84, 1988; Eriksen et al., Science 241: 84-86, 1988) as well as androgen (Colvard et al., Proc. Natl. Acad. Sci. 86: 854-857, 1989) receptors in osteoblasts could explain increased bone resorption observed after estrogen and androgen depletion.

While, in women, there is a rapid bone loss starting at menopause, bone loss in males can be recognized at about 65 years of age (Riggs et al., J. Clin. Invest. 67: 328-335, 1987). A more significant bone loss is seen in men at about 80 years of age, with the accompanying occurrence of hip, spine and wrist fractures. Several studies indicate that osteoporosis is a clinical manifestation of androgen deficiency in men (Baran et al., Calcif. Tissue Res. 26: 103-106, 1978; Odell and Swerdloff, West J. Med. 124: 446-475, 1976; Smith and Walker, Calif. Tissue Res. 22 (Suppl.): 225-228, 1976).

Therapy of postmenopausal women with nandrolone increased cortical bone mineral content (Clin. Orthop. 225: 273-277). Androgenic side-effects, however, were recorded in 50% of patients. Such data are of interest since while most therapies are limited to an arrest of bone loss, an increase in bone mass was found with the use of the anabolic steroid nandrolone. A similar

SUBSTITUTE SHEET

- 5 -

stimulation of bone formation by androgens has been suggested in a hypogonadal male (Baran et al., *Calcif. Tissue Res.* 26: 103, 1978).

5 The decline with age of serum levels of DHEA-S and DHEA has led to the intriguing possibility that low serum DHEA and DHEA-S levels could be associated with breast cancer and cardiovascular diseases. In fact, a series of studies have indicated that subnormal levels of DHEA are associated with a high risk of breast cancer (Bulbrook et al., *Lancet* 2: 395-398, 1971; Rose et al., *Eur. J. Cancer* 13: 43-47, 1977; Thijssen et al., *J. Steroid Biochem.* 6: 10 729-734, 1975; Wang et al., *Eur. J. Cancer* 10: 477-482, 1974; Gomes et al., *C.R. Acad. Sci. Paris* 306: 261-264, 1978; Brownsez et al., *Eur. J. Cancer* 8: 131-137, 1972). Women with breast cancer were found to have low urinary levels of androsterone and etiocholanolone, two metabolites of DHEA (Bulbrook et al., *Lancet* 2:1238-1240, 1962; Cameron et al., *Br. Med. J.* 4: 15 768-771, 1970). Bulbrook et al. (*Lancet* 2: 1235-1240, 1962) then reported that women with primary operable breast cancer had urinary levels of 11-deoxy-17-ketosteroids (derived mainly from DHEA-S and DHEA) lower than normal, thus suggesting that a low secretion rate of DHEA and DHEA-S could precede the development of breast cancer.

20 The main approaches for the treatment of already developed breast cancer are related to the inhibition of estrogen action and/or formation. The role of estrogens in promoting the growth of estrogen-sensitive breast cancer has been recognized (Lippman, *Semin. Oncol.* 10 (Suppl. 4): 11-19, 1983; Sledge and McGuire, *Cancer Res.* 38: 61-75, 1984; Witliff, *Cancer* 53: 630-643, 25 1984; Poulin and Labrie, *Cancer Res.* 46: 4933-4937, 1986).

DHEA (450 mg/kg, b.w., 3 times a week) markedly delayed the appearance of breast tumors in C3H mice which were genetically bred to develop breast 30 cancer (Schwartz, *Cancer Res.* 39: 1129-1132, 1979). Moreover, the risk of developing bladder cancer was found to be increased in men having lower serum DHEA levels (Gordon et al., *Cancer Res.* 51: 1366-1369, 1991).

Copending U.S. Patent Application No 07/785,890 filed November 4, 1991, 35 relates to a method of treatment of breast and endometrial cancer in susceptible warm-blooded animals which may include inhibition of ovarian

SUBSTITUTE SHEET

- 6 -

- hormonal secretion by surgical means (ovariectomy) or chemical means (use of an LHRH agonist, e.g. [D-Trp⁶, des-Gly-NH₂¹⁰]LHRH ethylamide, or antagonist) as part of a combination therapy. Antiestrogens, androgens, progestins, inhibitors of sex steroid formation (especially of 17 β -
- 5 hydroxysteroid dehydrogenase- or aromatase-catalyzed production of sex steroids), inhibitors of prolactin secretion and of growth hormone secretion and ACTH secretion are discussed. A counterpart thereof has been published under international publication number WO 90/10462.
- 10 Copending U.S. Patent Applications 07/724,532 and 07/900,817 filed on June 28, 1991 and June 24, 1992, respectively, relate to a method using low dose androgenic compounds for the prevention and treatment of breast cancer, endometrial cancer, osteoporosis and endometriosis. A counterpart of the 1992 application has been published as WO 93/00070.
- 15 Recent *in vitro* studies describe the relative antiproliferative activities of an androgen on the growth of the estrogen-sensitive human mammary carcinoma cell line ZR-75-1 (Poulin et al. "Androgens inhibit basal and estrogen-induced cell proliferation in the ZR-75-1 human breast cancer cell
- 20 line", Breast Cancer Res. Treatm. 12: 213-225, 1989). As mentioned above, Poulin et al. (Breast Cancer Res. Treatm. 12: 213-225, 1989) have found that the growth of ZR-75-1 human breast carcinoma cells is inhibited by androgens, the inhibitory effect of androgens being additive to that of an antiestrogen. The inhibitory effect of androgens on the growth of human
- 25 breast carcinoma cells ZR-75-1 has also been observed *in vivo* in nude mice (Dauvois and Labrie, Cancer Res. 51: 3131-3135, 1991).
- DHEA has been suggested to have beneficial effects in obesity, diabetes, atherosclerosis, chemically induced breast, skin and colon (prevention)
- 30 cancer, autoimmune diseases, fatigue, loss of muscle mass, connective tissue diseases, aging and longevity (Orentreich et al., J. Clin. Endocrinol. Metab. 59: 551-555, 1984; Regelson, Ann. N.Y. Acad. Sci. 521: 260-273, 1988; Gordon et al., Adv. Enzyme Regul. 26: 355-383, 1987; Schwartz, Adv. Cancer Res. 51: 391-423, 1988; Barrett-Connor et al., New Engl. J. Med.
- 35 315: 1519-1524, 1986).

SUBSTITUTE SHEET

- 7 -

In aged Sprague-Dawley rats, Schwartz (in Kent, Geriatrics 37: 157-160, 1982) has observed that body weight was reduced from 600 to 550 g by DHEA without affecting food intake. Schwartz (Cancer 39: 1129-1132, 1979) observed that C3H mice given DHEA (450 mg/kg, 3 times a week) gained significantly less weight and grew older than the control animals, had less body fat and were more active. The reduction in body weight was achieved without loss of appetite or food restriction. Furthermore, DHEA could prevent weight gain in animals bred to become obese in adulthood (in Kent, Geriatrics 37: 157-160, 1982).

DHEA in the diet has been shown to be a potent antihyperglycemic and antidiabetic agent in mice with inherited obesity-glucose intolerance syndrome (Coleman et al., Diabetes 33: 26-32, 1984).

DHEA reduced the incidence of atherosclerosis in cholesterol-fed rabbits (Gordon et al., J. Clin. Invest. 82: 712-720, 1988; Arad et al., Arteriosclerosis 9: 159-166, 1989). Moreover, high serum concentrations of DHEA-S have been reported to protect against death from cardiovascular diseases in men (Barrett-Connor et al., N. Engl. J. Med. 315: 1519-1524, 1986). Circulating levels of DHEA and DHEA-S have thus been found to be inversely correlated with mortality from cardiovascular disease (Barret-Connor et al., N. Engl. J. Med. 315: 1519-1524, 1986) and to decrease in parallel with the diminished immune competence (Thoman and Weigle, Adv. Immunol. 46: 221-222, 1989). A study in man has shown an inverse correlation between fetal serum DHEA-S and low density lipoprotein (LDL) levels (Parker et al., Science 208: 512, 1980).

In normal men, a placebo-controlled trial investigated the effect of daily oral administration of 1.6 g of DHEA for 28 days. Serum DHEA levels were increased 2.5- to 3.5-fold in the DHEA-treated group while total cholesterol and serum LDL cholesterol decreased by 7.1 and 7.5%, respectively (Nestler et al., J. Clin. Endocrinol. Metab. 66: 57-61, 1988). A decrease in body fat was found in 4 of the 5 men treated with DHEA for an average 31% decrease in percent of body fat with no change in total weight, thus suggesting a corresponding increase in muscle mass.

SUBSTITUTE SHEET

- 8 -

Drucker et al. (J. Clin. Endocrinol. Metab. 35, 48, 1972), Buster et al. (Am. J. Obstet. Gynecol. 166, 1163, 1992) and Welle et al. (J. Clin. Endocrinol. Metab., 71, 1259, 1990) have also administered DHEA orally to the human.

5

Obesity was found to be improved in the A^{VY} mutant mouse (Yen et al., Lipids 12: 409-413, 1977) and in the Zucker rat (Cleary and Zisk, Fed. Proc. 42: 536, 1983). DHEA-treated C3H mice had a younger appearance than controls (Schwartz, Cancer Res. 39: 1129-1132, 1979).

10

Brain concentrations of DHEA are 6.5 times higher than corresponding concentrations in plasma (Lacroix et al., J. Steroid Biochem. 28: 317-325, 1987). DHEA and DHEA-S improve memory in aging mice (Flood and Roberts, Brain Res. 448: 178-181, 1988). Serum DHEA-S concentrations in patients with Alzheimer disease have been found to be 48% lower on average than in age-matched controls (Sunderland et al., Lancet ii: 570, 1989). As mentioned above, DHEA administered chronically in the diet has been shown to increase longevity by delaying the development of some diseases in particular strains of animals.

15

U.S. Patent No 4,496,556 describes the use of DHEA or its derivatives to treat skin dryness by topical administration. Only local action on the sebaceous glands was described and no systemic action was observed.

20

U.S. Patent No 4,542,129 describes a topical composition for treating dry skin in a patient comprising the combination of DHEA and/or derivatives, a keratolytic agent and a non-toxic dermatologically acceptable vehicle.

Great Britain Patent No 1246639 describes preparation of esters of dehydroepiandrosterone for use as agents in the treatment of post and premenopause, tachycardia and headaches.

25

One problem facing the use of DHEA in humans is that high doses are required apparently because a large proportion of the compound is degraded in the liver before it reaches the blood stream after oral administration.

30

SUBSTITUTE SHEET

- 9 -

- It is known that the efficiency of delivery of some drugs can be improved by the use of certain pharmacologically inactive derivatives which are, by *in vivo* enzymatic or spontaneous reactions, transformed into the active drugs (see generally H. Bundgaard, Design and application of prodrugs. In A textbook of
- 5 Drug Design and Development. Edited by P. Krogsgaard-Larsen and H. Bundgaard. Harwood Academic Publishers GmFH, Chur, Switzerland, 1991, pp. 113-191). For example, Druzgala et al., J. Steroid Biochem. Molec. Biol. 38, 149-154, 1991, describes prodrugs of glucocorticoids. Bodor et al. in
- 10 U.S. Patent Appln. No 4,213,978 and in German Patent Application Publication No DE 29 48 733 disclose the use of thiazolidine derivatives of progesterone as topical drugs. Percutaneous absorption of prodrug derivatives of estrogens and progestins are reported by Friend DR in Critical Reviews in Therapeutic Drug Carrier Systems, vol. 7 (2), pp. 149-186, 1990. Information about percutaneous absorption of drugs can also be found in
- 15 Skin Permeability (H. Schaefer, A. Zesch and G. Stuttgen, eds), Springer-Verlag, Berlin, Heidelberg, New York, 1982, pp. 896.

- Currently, low dose estrogen therapy is the standard approach used in perimenopausal and menopausal women to relieve vasomotor symptoms,
- 20 urogenital atrophy, osteoporosis and other symptoms and signs associated with menopause (for review, see Edman, C.D., Estrogen Replacement Therapy. In: The menopause, Springer-Verlag, New York, (edited by H.J. Buchsbaum), pp. 77-84, 1983). Detailed information about the menopause and its therapy can be found in other chapters of this book. Obviously, such
- 25 approaches limited to estrogen replacement therapy, associated or not with progestins, do not reproduce the equilibrium between estrogens and androgens that occurs naturally from the transformation of DHEA into its active metabolites in various target tissues.

- 30 Some esters of DHEA at position 3 are already described in the literature (Riva et al., J. Org. Chem. 54: 3161-4, 1989; Parish and Chistrakorn, Synth. Commun. 15: 393-9, 1985; Rom Patent No RO 66924B; Jarosz and Zamojski, Tetrahedron 38:1453-6, 1982; Heublin et al., Z. Chem. 22: 178, 1982; German Patent Application No DE 2534911; Khaidem et al., Indian J. Chem.
- 35 Sect. B, 27B: 850-1, 1988; Pettit et al., J. Org. Chem. 52: 3573-8, 1987; Hanson and Reese, J. Chem. Soc. Perkin Trans. 1: 647-649, 1985);

SUBSTITUTE SHEET

- 10 -

European Patent Application No 84-105741; Heublein et al., Acta Polym., 35: 673-7, 1984; Seevers et al., J. Med. Chem., 25: 1500-3, 1982; Yamashita and Kurosawa, Agric. Biol. Chem., 39: 2243-4, 1975; Japan Patent Application JP 50005372; Pohlmann et al., Mol. Cryst. Liquid Cryst. 13: 243-54, 1971.

5

Alkanesulfonates of DHEA are described as inhibitors of glucose-6-phosphate dehydrogenase activity in J. Pharm. Sci. 73: 1643-5, 1984.

10 In Britain Patent Application No GB 1246639 and S. Africa Patent Application No ZA 6806112 discloses DHEA esters for the treatment respectively of post and premenopause tachycardia and headaches and climateric complaints.

Leszczynski et al., in Biochem. Biophys. Acta, 1014: 90-7, 1989; idem: 1083: 18-28, 1991, have reported esterification of DHEA by blood plasma and Katz
15 et al., in MCF-7 cell line in J. Steroid Biochem, 26: 687-92, 1987.

Ethyl carbonate of DHEA is reported by Weisz and Agocs in Arch. Pharm. (Weinheim, Ger), 319: 952-3, 1986.

20 Some halogeno esters of DHEA are described by Challis and Heap in J. Chromatogr. 50: 228-238, 1970 and by Pinelly and Nair in J. Chromatogr. 43: 223-228, 1969.

25 Although DHEA has been suggested as involved in various biological functions, as discussed above, its pharmaceutical use as a therapeutic or prophylactic agent has been relatively limited. Its role in preventing, reducing or even reversing progress of certain diseases has not heretofore been fully understood. The present invention now discloses a number of new pharmaceutical uses of DHEA and of DHEA-S (or compounds convertible *in*
30 *vivo* to either). The invention also provides improved methods of administering these agents which can overcome the disadvantages associated, for example, with oral administration.

SUBSTITUTE SHEET

- 11 -

SUMMARY OF THE INVENTION

It is an object of the present invention to provide simple and efficient methods for systemic delivery of sex steroid precursors such as dehydroepiandrosterone, dehydroepiandrosterone sulfate and analogs which
5 are converted *in vivo* to either DHEA or DHEA-S.

It is another object of the invention to provide novel methods of preventing and/or treating treatments of menopause, vaginal atrophy, hypogonadism, diminished libido, osteoporosis, loss of skin thickness and cellularity (skin
10 atrophy), urinary incontinence, ovarian cancer and uterine cancer.

It is another object of the invention to provide kits and pharmaceutical compositions for use in accordance with the invention.

15 It is another object to provide novel contraceptive method.

It is another object of the invention to provide novel sex steroid precursors and pharmaceutical compositions thereof.

20 In one aspect, the invention provides a method for treating menopause comprising administering to a patient in need of such treatment an effective amount of at least one sex steroid precursor selected from the group consisting of dehydroepiandrosterone, dehydroepiandrosterone sulphate, and compounds converted *in vivo* to either of the foregoing, in combination
25 with an effective amount of an estrogen, a progestin or both.

In another aspect, the invention provides a pharmaceutical composition for the treatment of menopause and other indications discussed herein comprising at least one sex steroid precursor selected from the group
30 consisting of dehydroepiandrosterone, dehydroepiandrosterone sulphate, and compounds converted *in vivo* to either of the foregoing, and further comprising an estrogen or a progestin or both. One preferred combination is precursor and estrogen. Another is precursor and progestin. In some preferred embodiments, precursor, progestin and estrogen are used in
35 combination. Estrogen is disfavored for male patients.

SUBSTITUTE SHEET

- 12 -

In another aspect, the invention provides a kit for the treatment of menopause having a first container which includes at least one sex steroid precursor selected from the group consisting of dehydroepiandrosterone, 5 dehydroepiandrosterone sulphate, and compounds converted *in vivo* to either of the foregoing, and at least one additional container having either a progestin, an estrogen, or both. One preferred kit includes three containers having a precursor, an estrogen and a progestin, respectively, in separate containers.

10

However, 2 or more of the active ingredients may, if desired, be in a single container. A pharmaceutical carrier or diluent may also be provided in one or more of the containers and may include preservatives and many other additives known in the art.

15

In another aspect, the invention provides a method of treating vaginal atrophy, hypogonadism, diminished libido, reduced skin thickness and cellularity comprising administering to a patient in need of such treatment an effective amount of a sex steroid precursor selected from the group consisting 20 of dehydroepiandrosterone, dehydroepiandrosterone sulphate, and compounds converted *in vivo* to either of the foregoing.

In another aspect, the invention provides a method for the prevention or treatment of osteoporosis comprising administering to a patient in need of 25 such prevention or treatment a therapeutically effective amount, with or without a pharmaceutical diluent or carrier, of a sex steroid precursor selected from the group consisting of DHEA, DHEA-S and compounds that are converted *in vivo* to either.

30 In another aspect, the invention provides a method for the prevention or treatment of urinary incontinence comprising administering to a patient in need of such prevention or treatment, with or without additional pharmaceutical diluent or carrier, a therapeutically effective amount of a sex steroid precursor selected from the group consisting of DHEA, DHEA-S and 35 compounds converted *in vivo* to either.

SUBSTITUTE SHEET

- 13 -

In another aspect, the invention provides a method of contraception comprising administering to a female in need of contraception, with or without additional pharmaceutical diluent or carrier, an effective amount of a sex steroid precursor selected from the group consisting of DHEA, DHEA-S and compounds converted *in vivo* to either. In preferred embodiments, an estrogen and/or progestin are further administered as part of a combination method of contraception together with the precursor.

In another aspect, the invention provides the pharmaceutical composition of a sex steroid precursor selected from the group consisting of DHEA, DHEA-S and compounds converted *in vivo* to either, and a supplemental agent selected from the group consisting of an estrogen and a progestin. In certain embodiments, estrogen and progestin are both included. Preferably, a pharmaceutical diluent or carrier is also added in some embodiments.

It is also possible to make kits in accordance with the invention which provide at least two separate containers, one of which includes the sex steroid precursor, and another of which includes either an estrogen or a progestin or both. In certain embodiments, three separate containers may be provided wherein one container has at least the sex steroid precursor therein, another container has at least the estrogen therein, and another container has at least the progestin therein. All indications discussed herein as responding to the sex steroid precursors (DHEA, DHEA-S and compounds converted *in vivo* to either) may, in certain embodiments, further respond to administering estrogen and/or progestin in combination with the precursor (with the exception of treatment in men where an estrogen would be disfavored). Thus, the kits and pharmaceutical compositions should provide the combination of the foregoing agents appropriate to the particular indication for which they will be used, and the combination therapy chosen.

In another embodiment, the invention provides a method for prevention of ovarian cancer comprising administering to a female patient in need of such prevention, with or without additional pharmaceutical diluent or carrier, a therapeutically effective amount of a sex steroid precursor selected from the group consisting of DHEA, DHEA-S and a compound converted *in vivo* to either.

SUBSTITUTE SHEET

- 14 -

In another embodiment, the invention provides a method for prevention of uterine cancer comprising administering to a female patient in need of such prevention, with or without additional pharmaceutical diluent or carrier, a therapeutically effective amount of a sex steroid precursor selected from the group consisting of DHEA, DHEA-S and a compound converted *in vivo* to either.

In another aspect, the invention provides a therapeutic method of treating reduced or imbalanced concentrations of sex steroids comprising applying an effective amount of a pharmaceutical composition for percutaneous or transmucosal delivery to an outer surface of skin or mucosa of a patient in need of such treatment, said pharmaceutical composition comprising a carrier having dissolved therein at least one sex steroid precursor selected from the group consisting of dehydroepiandrosterone, dehydroepiandrosterone sulphate, and compounds converted *in vivo* to either of the foregoing, said precursor being present at a concentration of at least 7% by weight relative to the total pharmaceutical composition, and said carrier being compatible with skin or mucosa and allowing penetration of said precursor through said skin or mucosa, said carrier having sufficient viscosity to maintain said precursor on a localized area of skin or mucosa, without running or evaporating, for a time period sufficient to permit substantial penetration of said precursor through said localized area of said skin or mucosa. The foregoing method is useful in treating and/or preventing the conditions discussed above, menopausal symptoms and other conditions which respond to replenishment of diminished DHEA levels, including but not limited to obesity, cardiovascular disease, atherosclerosis, breast cancer, endometrial cancer, loss of muscle mass, diabetes, fatigue, connective tissue diseases and memory loss.


In another aspect, the invention provides a pharmaceutical composition for percutaneous or transmucosal delivery of said pharmaceutical composition comprising a carrier having dissolved therein at least one sex steroid precursor selected from the group consisting of dehydroepiandrosterone, dehydroepiandrosterone sulphate, and compounds converted *in vivo* to either of the foregoing, said precursor being present at a concentration of at least

SUBSTITUTE SHEET

- 15 -

7% by weight relative to the total pharmaceutical composition, and said carrier being compatible with skin or mucosa and allowing penetration of said precursor through said skin or mucosa, said carrier having sufficient viscosity to maintain said precursor on a localized area of skin or mucosa, without
 5 running or evaporating, for a time period sufficient to permit substantial penetration of said precursor through said localized area of said skin or mucosa.

In another aspect, the invention provides novel compounds (and
 10 pharmaceutical compositions containing them) of the formulas set forth below with substituent definitions set forth below:



15 wherein X is selected from the group consisting of H, $R - \overset{\overset{O}{\parallel}}{C} -$, RCO_2CHR^a - and R^bSO_2 -

R being selected from the group consisting of hydrogen, straight- or branched-alkyl, straight- or branched-alkenyl, straight- or branched-alkynyl,
 20 aryl, furyl, straight- or branched-alkoxy, straight- or branched-alkenyloxy, straight- or branched-alkynyloxy, aryloxy, furyloxy and halogeno analogs of the foregoing


R^a being hydrogen or (C₁-C₆) alkyl; and
 25

R^b being selected from the group consisting of hydroxyl (or salts thereof), methyl, phenyl and p-toluy;

SUBSTITUTE SHEET


- 16 -

wherein Y is a divalent ;substituted or unsubstituted moiety of the formula:




(Z being an oxygen or sulfur atom), and wherein Y and the carbon atom to which it is bonded together form a closed saturated 5-membered ring.

5



wherein R^c is selected from the group consisting of C_3 - C_{20} straight- or branched-alkyl, C_3 - C_{20} straight- or branched-alkenyl, C_3 - C_{20} straight- or branched-alkynyl, aryl and halo-substituted analogs of the foregoing.

10



R^d being selected from the group consisting of hydrogen, straight- or branched-alkyl, straight- or branched-alkenyl, straight- or branched-alkynyl, aryl, furyl, straight- or branched-alkoxy, straight- or branched-alkenyloxy, straight- or branched-alkynyloxy, aryloxy, furyloxy and halogeno analogs of the foregoing.


15

wherein R_a is hydrogen or (C_1 - C_6) alkyl.

20

SUBSTITUTE SHEET

- 17 -



wherein R³ is hydroxy or sulfate.

- 5 wherein R^e is selected from the group consisting of hydrogen, benzyl, aryl, straight- or branched-alkyl, straight- or branched-alkenyl and straight- or branched-alkynyl.

wherein Z is oxygen or sulfur.

10

In one embodiment, a method is provided to compensate for the marked decrease in the secretion of the sex steroid precursors DHEA and DHEA-S by the adrenals during aging comprising administering DHEA, DHEA-S or analogs converted *in vivo* thereto in amounts which compensate for the consequences of decreased DHEA and DHEA-S secretion by the aging adrenals without exerting unwanted side effects.

15

It is believed that the methods of the invention are suitable for both prophylactic and therapeutic use. The serum concentrations, kits, and compositions discussed herein are equally useful toward either objective.

20

In another aspect, the invention provides a transdermal device comprising;

(a) a surface adapted for contact with human skin;

(b) a means of maintaining said surface on a localized area of skin to which said device is applied;

25

(c) a storage member in fluid communication with said surface, said member containing a pharmaceutical composition comprising a carrier and an active ingredient selected from the group consisting of dehydroepiandrosterone, dehydroepiandrosterone sulphate and analogues which are converted *in vivo*

30

to either; and

SUBSTITUTE SHEET.

- 18 -

(d) a means for conducting said pharmaceutical composition from said storage member through said surface and into contact with said localized area of skin.

- 5 For all indications for which DHEA is recommended herein in accordance with the present invention, it is to be understood that any sex steroid precursor selected from the group consisting of DHEA, DHEA-S and compounds converted *in vivo* to either (e.g. prodrug forms of DHEA or DHEA-S) may be used. All will result in increased serum levels of DHEA. Because
- 10 DHEA-S is a natural precursor of DHEA, DHEA-S (as well as prodrugs to either of DHEA or DHEA-S) may be used in place of DHEA for the various indications discussed herein for using DHEA. The result of administering these prodrugs or of administering DHEA-S is desirably increased levels of DHEA.

15

DESCRIPTION OF THE DRAWINGS

- 20 Figures 1-3 show the increase in bone density of rats after nine months of treatment with DHEA (EM-760). Figure 1 shows total bone mass density of both intact and ovariectomized rats treated for 9 months with DHEA versus an untreated control group of rats. Figure 2 shows the same comparison measuring lumber spine mass density, and Figure 3 shows the same comparison measuring femoral bone mass density. In each case, except of
- 25 course for the intact control group, measurements were taken nine months after ovariectomy and/or percutaneous DHEA administration. These figures illustrate the value of the present invention in the treatment or prevention of osteoporosis.
- 30 Figures 4 and 5 illustrate the effectiveness of DHEA as a contraceptive, or for therapy or prevention of ovarian or uterine cancer. Figure 4 illustrates the reduction in rat ovarian weight achieved after six months treatment with twice daily percutaneous administration of DHEA at a dose of 30 mg in 0.5 ml 50% ethanol-50% propyleneglycol on the dorsal skin area covering about two
- 35 square centimeters thereof.

SUBSTITUTE SHEET

- 19 -

Figure 5 illustrates histology of the ovaries of intact control rats (Fig. 5A) and intact rats treated with DHEA at a dose of 30 mg twice daily in a solution of 50% ethanol - 50% propylene glycol applied on an area of 2 cm² of dorsal skin (Fig. 5B). Figure 5B shows a marked decrease in the number of tertiary and secondary follicles (F) and absence of corpora lutea (CL); interstitial cells (IC). Magnification x 250. Note the atrophy of the interstitial gland in the treated rat (Fig. 5C) in comparison with the intact (Fig. 5D) at magnification x 500.

Figure 6 illustrates the effect on vaginal atrophy of one, three or six months of treatment with DHEA administered at a dose of 30 mg twice daily in a solution of 50% ethanol - 50% propylene glycol on an area of 2 cm² of dorsal skin in the ovariectomized rat. Atrophic vaginal epithelium is shown after 1 (Fig. 6A), 3 (Fig. 6B), and 6 months following castration (Fig. 6C). Reversal of vaginal atrophy in rats treated with DHEA as shown in Figs 6D, 6E and 6F illustrating the vaginal epithelium after DHEA treatment for 1, 3, and 6 months, respectively. Magnification x 200. Thus, histopathologic examination showed proliferation and murification of the vaginal epithelium and reversal of vaginal atrophy in the rats treated with DHEA.

Figure 7 shows histology of the skin of male rats after 3 and 6 months of treatment with DHEA given by topical application in an area of 2x2 cm on dorsal skin. Slight to moderate hypertrophy and hyperplasia of sebaceous glands in dorsal (b) and (c); as well as in ventral (e) and (f) skin of intact treated rats for 3 months (b) and (e) and 6 months (c) and (f) compare with intact controls (a) and (d). Distention of ducts (D).

Figure 8 shows histology of the skin of male rats after 3 and 6 months of treatment with DHEA given by topical application in an area of 2 x 2 cm on dorsal skin. Slight to moderate hyperplasia of sebaceous glands in dorsal (b) and (c); as well as in ventral (e) and (f) skin of castrated rats treated for 3 months (b) and (e) and 6 months (c) and (f). Compare with castrated controls (a) and (d). Note the distention of ducts (D) and increase in the number of the acini (A) per hair follicle. Magnification x 100.

SUBSTITUTE SHEET

- 20 -

Figure 9. Effect of treatment with DHEA on dorsal skin dermis (area of topical application of DHEA) in male castrated rats after 3 (c) and 6 months (d) and (f). Castrated untreated rats were used as controls (a), (b) and (e). Increase in thickness and cellularity of dermis in treated rats (c), (d) and (f). a, b, c, d: 5 magnification x 100; e, f. magnification x 500.

Figures 7 - 9 above illustrate that topical application of DHEA on the dorsal skin leads to an increased thickness and cellularity of the dermis, an effect which should prevent or treat skin atrophy (a condition marked *inter alia* by 10 loss of collagen connective tissue). The effects of DHEA are also visible on the size of the sebaceous glands which are believed to be stimulated by the androgenic compounds produced from DHEA as a precursor.

As can be seen in Figs. 7 and 8, topical application of DHEA (30 mg in 50% 15 ethanol/50% propylene glycol) for 3 or 6 months on an area of 2 x 2 cm on dorsal skin leads to a slight to moderate hypertrophy and hyperplasia of the sebaceous glands in both the dorsal and ventral skin areas, thus indicating a generalized effect of this treatment on the skin. This effect is seen in both intact and castrated animals where the marked atrophy following castration is 20 completely prevented by DHEA treatment. Such a decrease in the size of the sebaceous glands after castration can be compared to the atrophy occurring during aging.

As illustrated in Fig. 9, the effect of DHEA is accompanied by a marked 25 increase in the thickness and cellularity of the dermis in both intact and castrated animals. Since collagen is an important component of the skin dermis, the present data indicate that the correction of skin atrophy could be achieved, at least in part, by increased formation of collagenous tissue.

30 Regarding the effectiveness of DHEA as a contraceptive, topical administration of DHEA (30 mg) in 50% ethanol - 50% propylene glycol twice daily on an area of 2 cm² on the dorsal skin of intact rats led to histopathological changes that demonstrate inhibition of ovulation. The most 35 important changes observed were an atrophy of the ovaries (Fig. 4) and histopathological changes (Fig. 5) indicating an absence of ovulatory cycles. There was a marked decrease in the number of secondary and tertiary

SUBSTITUTE SHEET.

- 21 -

follicles and an absence of corpora lutea, thus showing the absence of ovulation. These histopathological signs of absence of ovulation in the ovaries were also accompanied by an arrest of the cyclic histopathological changes normally seen in the endometrium during the estrous cycle in the rat. These histopathological changes show that treatment with DHEA exerts contraceptive action.

The invention is further explained in the following non-limiting description of preferred embodiments.

DETAILED DESCRIPTION OF THE PREFERRED EMBODIMENTS

To select patients who may benefit from the treatments described herein, the serum levels of DHEA and its metabolites can be measured as described by Belanger et al., in Steroid Formation, Degradation and Action in Peripheral, Normal and Neoplastic Tissues (H Bradlow, L Castagnetta, S d'Aquino, L Gogliotti, eds) Ann. N.Y. Acad. Sci. 586: 93-100, 1990; Haning et al., J. Clin. Endocrinol. Metab. 72: 1088, 1991. See also Labrie et al., Endocrinology 123, 1412-1417, 1988. Serum IGF-1 levels can be measured as described (Furlanetto et al., J. Clin. Invest. 60: 648, 1977). In accordance with the invention, once the DHEA deficiency is determined, DHEA or its analogues is preferably administered at a dosage sufficient to cause and maintain serum DHEA concentration between 4 and 10 micrograms per liter, especially between 4 and 7 micrograms per liter. Higher concentrations are desirable in certain indications as discussed below.

In some preferred embodiments, serum concentration is between 5 and 7 or between 6 and 7 micrograms per liter. However, for purposes of contraception or for prevention of ovarian or uterine cancer, concentration up to 13 micrograms per liter (e.g. between 7 and 13) are preferred. Preferred dosages discussed herein may be increased as appropriate to achieve these higher serum concentrations, e.g. by a factor of about 30% with variations for individual patient response as monitored by the attending clinician. When DHEA is administered by the preferred percutaneous or transmucosal

SUBSTITUTE SHEET

- 22 -

technique, it has been found that DHEA is very efficiently absorbed into the blood to raise serum levels. For example, when a Glaxal cream, containing Glaxal base as a carrier (available from Glaxal Canada Limited) and 10% DHEA by weight of the total composition, is applied twice daily to a 100 square centimeter surface of the abdominal area in an amount providing 100 milligrams of active ingredient (e.g. DHEA), a typical patient is likely to respond with an increase of serum DHEA concentration of about 0.7 micrograms per liter per 50 kg of body weight. The delivered dosage may be raised or lowered in known manners by altering the location to which the lotion or ointment is applied, by altering the size of the surface area to which it is supplied, by altering the concentration of the active ingredient, or by altering the carrier. For example, increasing the surface area will normally increase the dosage of active ingredient delivered if the concentration of active ingredient remains constant. In the same manner, dosage delivered increases with increased concentration of active ingredient in the delivery base, and decreases with decreased concentration. Dosage delivered into the bloodstream also varies in a known manner with respect to the body region at which the transdermal penetration system is applied to the skin. Changing the carrier can also alter the delivered dosage in known ways. Preferably, serum DHEA concentration is measured before treatment begins, and a dosage is selected to quickly raise serum DHEA concentration to the preferred target range between 4 and 10 micrograms per liter or 7 to 13 micrograms per liter for the higher dosage indications discussed above. Subsequently, the patient is monitored both symptomatologically and by DHEA concentration to verify that the desired serum concentration target and symptomatic relief have been obtained. DHEA is then maintained at a constant concentration in the circulation. For a typical post-menopausal patient, for example, this dosage is the equivalent of application of 400 mg of the active precursor, as part of a 10 percent composition in Glaxal, to a 400 square centimeter area of the abdomen 2 times daily per 50 kg of body weight. If oral administration is chosen, 800 mg should be administered twice daily per 50 kg of body weight.

In accordance with the invention, DHEA, DHEA-S and/or compounds converted to either *in vivo* are utilized for the treatment and/or prevention of menopausal symptoms, vaginal atrophy, atrophy of the skin, hypogonadism,

SUBSTITUTE SHEET

- 23 -

diminished libido, osteoporosis, urinary incontinence, ovarian cancer or uterine cancer. Additionally, other conditions related to decreased secretion of DHEA by the adrenals during aging and which respond to DHEA therapy can be treated more efficiently with transdermally delivered DHEA, DHEA-S (or analogues) in accordance with the invention. Conditions expected to respond to the treatments herein may be diagnosed in conventional ways. For example, the appearance of breast cancer is usually detected by self breast examination, clinical breast examination by the physician and/or mammography. Endometrial cancer, on the other hand, is usually diagnosed by the PAP smear and/or endometrial biopsy. Both cancers can be diagnosed and evaluated by standard physical methods well known to those skilled in the art, e.g. bone scan, chest X-Ray, skeletal survey, ultrasonography of the liver and liver scan (if needed), CAT scan, MRI and physical examination.

The first manifestations of menopause are usually hot flashes. Further characterization of menopause can be determined in accordance with known techniques. See for Example, The Menopause (Herbert J. Buchsbaum, ed), Springer Verlag, New York (1983), pp. 222. Vaginal atrophy is often indicated by dyspareunia and vaginal infections. Vaginal atrophy, hypogonadism and diminished libido are all characterized in well-known ways. For the above-indicated diseases, see, for example, Korenman, Stanley G, "Sexual Dysfunctions" in Williams Textbook of Endocrinology (Jean D. Wilson and Daniel W. Foster, Eds.), WB Saunders Co, Philadelphia, pp. 1033-1048, 1992.

Bone density, on the other hand, can be measured by standard methods well known to those skilled in the art, e.g. QDR (Quantitative Digital Radiography), dual photon absorptiometry and computerized tomography. Plasma and urinary calcium and phosphate levels, plasma alkaline phosphatase, calcitonin and parathormone concentrations, as well as urinary hydroxyproline and calcium/creatinine ratios are useful parameters of bone formation and resorption.

SUBSTITUTE SHEET

- 24 -

Loss of collagen or connective tissues in the skin often accompanies aging, especially in persons over 50 years of age. It may be evidenced by wrinkling of the skin and/or low elasticity.

- 5 Osteoporosis or otherwise insufficient bone mass, and other diseases treatable by activating the androgen receptor may be treated in accordance with the present invention or prophylactically prevented in accordance herewith. The present invention can aid in the prevention of breast, ovarian or endometrial cancer.

10

The normal range of body weight is well known to those skilled in the art, while cholesterol and lipoproteins are routinely measured by standard techniques (Nestler et al. J. Clin. Endocrinol. Metab. 66: 57-61, 1988 for references).

15

Skin status can be assessed by visual inspection, palpation and, with more precision, by punch biopsy and standard histological examination.

20

The traditional main mechanism for female contraception in the prior art related to administering an estrogen, which at increased circulating levels, reduced LHRH secretion from the hypothalamus which, in turn, decreased LH secretion from the pituitary. The resultant reduction in LH secretion decreased ovarian function, and in particular ovulation. Addition of a progestin controlled the growth of the endometrium and transformed the vaginal and cervical secretions into an unfavorable environment for sperm capacitation and fertility.

25

In the present invention, DHEA is used instead of estrogen (although estrogen may also be added in certain embodiments discussed below). In accordance with the invention, DHEA provides estrogen for contraception while simultaneously and desirably providing increased levels of androgens which will contribute to contraception since androgens inhibit LHRH and LH secretion. These androgens can, especially in women at perimenopause (as well as in postmenopausal women when contraception is no longer required), provide much needed stimulation of bone formation and resistance to bone

30

SUBSTITUTE SHEET

- 25 -

loss. The estrogens produced from the administered DHEA also contribute to reducing bone loss. As with other uses discussed herein, use of DHEA instead of a sex steroid (here estrogen) avoids externally administering relatively high doses of estrogens and this avoids giving such estrogens extensive access to all tissues, many of which do not require estrogens. By substituting DHEA, estrogens are instead produced by natural processes in the same tissues where estrogens and androgens are needed and that normally convert DHEA to estrogens and androgens. The relative proportions of estrogen and androgen also remain substantially at natural levels in each specific tissue.

As with other uses of DHEA discussed herein, DHEA-S or prodrugs of DHEA or DHEA-S may be substituted instead or in addition to DHEA. Because ovarian function is diminished by the contraceptive technique described herein, ovarian production of estrogen and progesterone is decreased. Thus administering a progestin (e.g. medroxyprogesterone acetate, megestrol acetate, norethynodrel, L-norgestrel) to prevent endometrial hypertrophy as part of the contraceptive method is preferred. Androgenic progestins are preferred. The progestin may be administered in a pharmaceutical composition that includes the DHEA or separately. In certain embodiments, the progestin may be administered intermittently every month for 12-14 days, or 12-14 days every few months (e.g. every 2-4 months) or continuously. Progestin dosage may be in the range utilized in the prior art but is preferably lower for reasons explained below.

Estrogen may also be added to the contraceptive therapy because of decreased estrogen production in the ovaries. However, DHEA itself is converted to estrogen in many tissues and externally added estrogen can be minimized during practice of the present invention. Preferred dosage of added estrogen, when used in the contraceptive method is an amount effective to achieve between 100 and 200 nanograms estradiol per liter or equivalent. Preferably the ratio of added estradiol to DHEA (w/w) will range from 1,000 to 25,000 preferably 2,000 to 15,000 and especially 3,000 to 12,000. As with added progestin, added estrogen may be administered as part of a pharmaceutical composition that includes the DHEA (or, where used, the DHEA-S or prodrug) or separately. In some embodiments, DHEA,

SUBSTITUTE SHEET

- 26 -

progestin and estrogen are all administered, together or separately, as part of a combination therapy. A combination therapy results whenever a regimen of treatment elevates blood levels of each active agent simultaneously. This simply requires that the active agents be administered sufficiently close in
5 time that elevated blood levels of these agents were concurrent.

The use of combination contraceptives containing estrogens and progestins has not been shown to reduce the risk of breast cancer (Romiev et al., 1990, Cancer 66: 2253-63). These data are consistent with a known mitogenic effect of both estrogen and progesterone on breast cell epithelial
10 proliferation, thus explaining a peak of cell proliferation at mid-luteal phase (Masters et al., J. Natl. Cancer Inst. 1977, 58: 1263-65; Anderson et al., 1982, Br. J. Cancer 46: 376-82). In fact, total breast cell proliferation rate in premenopausal women using contraceptives is not different from that of untreated cycling women (Potter et al., 1988, Br. J. Cancer 58: 163-170;
15 Going et al., 1988, Am. J. Pathol. 130: 193-204).

Applicant has recently discovered that DHEA is transformed preferentially into androgens relative to estrogens. Thus, in accordance with the invention, this prohormone is used as a contraceptive instead of the current pill in order to desirably reduce breast cell proliferation. Androgens do in fact exert
20 inhibitory effects on breast cell proliferation by two mechanisms, namely a direct inhibitory effect in breast cells, and an inhibitory effect on gonadotropin secretion at the hypothalamo-pituitary level, thus resulting in decreased ovarian activity (e.g. less estrogen secretion and thus less estrogen-induced breast cell growth).

25 In addition to its beneficial effect on breast cells, the DHEA-induced decrease in gonadotropin secretion reduces ovarian activity (Figs. 4 and 5) and should thus help prevent ovarian cancer as well as uterine cancer. The increased gonadotropin secretion which precedes menopause will thus be ameliorated by DHEA administration. In accordance with the invention, DHEA may
30 therefore be used prophylactically against development of ovarian or uterine cancer in patients at high risk of these cancers even when contraception is not a primary objective.

SUBSTITUTE SHEET

- 27 -

- Prior art oral contraceptives add important amounts of the sex steroids estrogens and progestins. The present approach, however, is based on the tissue-specific formation and action of predominant androgens synthesized from DHEA. In fact, when peripheral tissues are exposed to the precursor steroid DHEA, these tissues make androgens predominantly as well as some estrogens at the site of action, the balance between the two categories of sex steroids being more physiological. DHEA will also block gonadotropin secretion and thus protect the ovary from hyperstimulation by high levels of gonadotropins preceding and accompanying menopause. The large proportion of the inhibition of gonadotropin secretion will be exerted by the DHEA-derived natural androgens, thus reducing to a minimum the needs for estrogens and progestin. This is important because many progestins including norethindrone and norethynodrel possess strong estrogenic activity (Poulin et al., Breast Cancer Res. Treat. 13: 265-276, 1989).
- Applicant has now discovered that the precursor steroid DHEA is converted to androgens (and estrogens) in osteoblasts (bone-forming cells). This discovery shows that, in accordance with the invention, DHEA may now be used instead of androgen and estrogen in the treatment or prevention of osteoporosis. Androgens produced in the bone (by conversion of the administered DHEA) stimulate bone formation and reduce bone loss, while estrogens produced from the administered DHEA also contribute to reducing bone loss. Significant side effects of traditional androgen therapy are thus avoided. For example, externally administered androgens of the prior art have access to many tissues that neither produce nor require androgens, thus causing side effects and disturbing the physiological balance of sex steroids in those tissues. By substituting DHEA (or prodrugs or DHEA-S if desired) in accordance with the invention, the DHEA is transformed to androgens only by natural mechanisms in tissues that normally perform that transformation according to their local needs. The relative ratio of androgens and estrogens produced from the DHEA is also a substantially normal ratio instead of being an abnormally elevated ratio of one type of sex steroid when that type alone is used.

In preferred embodiments, DHEA is administered for prevention or treatment of osteoporosis at a dosage sufficient to maintain substantially normal serum

SUBSTITUTE SHEET

- 28 -

- concentration for young adults, approximately 4-10 micrograms per liter, or in some embodiments 4-7 micrograms per liter, e.g. between 5 and 7 or between 6 and 7 micrograms per liter. These are also desirable levels for the other DHEA-responsive indications discussed herein, except for
- 5 contraception and prevention of ovarian and uterine cancer, where in some embodiments preferred dosage could be increased up to 13 micrograms per liter as discussed in connection with those particular indications in order to further inhibit LH secretion by the anterior pituitary gland.
- 10 In one preferred treatment for menopause, the invention seeks to simultaneously maintain blood levels of estrogen and a sex steroid precursor (e.g., DHEA or DHEA-S) within normal pre-menopausal parameters. The body converts DHEA-S to DHEA in most peripheral tissues. Without
- 15 intending to be bound by theory, it is believed that maintenance of appropriate precursor levels will better enable natural enzymes, such as 17 β -hydroxysteroid dehydrogenase, 3 β -hydroxysteroidhydrogenase, aromatase and 5 α -reductase to regulate production of androgens and estrogens and to maintain them in a manner more closely resembling their relative levels prevailing prior to menopause. Hence, the invention contemplates that not
- 20 only estrogens but also androgens and precursors will be kept in better balance. In fact, all target tissues possess the enzymatic machinery necessary to synthesize androgen and/or estrogens according to local control and need (Labrie, Mol. Cell. Endocrinol. 78, C113-C118, 1991).
- 25 The estrogen and precursor (e.g., DHEA) when administered together in accordance with the invention may be administered simultaneously or separately. Indeed, the second active ingredient (precursor) may be added to an existing estrogen therapy in order to achieve the combination therapy of the invention.
- 30
- It is necessary only that both the estrogen and precursor be administered in a manner and at a dosage sufficient to allow blood serum concentration of each to obtain desired levels. In accordance with the combination therapy of the invention, concentration of the precursor is maintained within desired

SUBSTITUTE SHEET

- 29 -

parameters at the same time that estrogen concentration is maintained within desired parameters. Where estradiol is used, serum estradiol concentration should typically be maintained between 50 and 300 nanograms per liter, preferably between 100 and 200 nanograms per liter and most preferably
5 between 150 and 175 nanograms per liter. Where another estrogen is used, serum concentration may be varied in a known manner to account for the difference in estrogenic activity relative to estradiol and in order to achieve normal pre-menopausal estrogen levels. A lesser concentration is needed, for example, if Mestranol is used. Adequate serum estrogen levels can also
10 be assessed by disappearance of the symptoms of menopause. Serum concentration of the second compound of the combination therapy (e.g., DHEA) is typically maintained between 4 and 10 micrograms per liter, or in some embodiments between 5 and 7 micrograms per liter, or between 6 and 7 micrograms per liter.

15 The estrogen is preferably estradiol, but may be sodium estrone sulfate or another compound which acts as an estrogen receptor agonist. When administered separately, commercially available estrogen supplements may be used, e.g., PREMARIN available from Ayerst (St-Laurent, Québec,
20 Canada). One preferred precursor is DHEA, although DHEA-S and analogs discussed below are also especially effective for the reasons stated below. Where DHEA is used, for example, DHEA, pharmaceutical grade, is commercially available from SIGMA (St-Louis, Missouri, USA). For typical patients, the appropriate dosage of estrogen to achieve desired serum
25 concentrations is between 0.3 and 2.5 milligrams of PREMARIN per day per 50 kg of body weight when administered orally. In certain embodiments of the invention, the estrogen may be 17 β -estradiol administered percutaneously in a patch which is available from CIBA under the name ESTRADERM wherein the daily dose is between 0.05 and 0.2 milligrams per
30 day per 50 kg of body weight. For typical patients, the appropriate dosage of the sex steroid precursor DHEA to achieve desired serum concentration of the precursor is between 0.25 and 2.5 grams per day per 50 kg of body weight when administered orally. Other sex steroid precursors will be administered at a dosage that depends on their *in vivo* conversion rate to
35 DHEA. The precursor may also be administered transdermally, as described

SUBSTITUTE SHEET

- 30 -

in more detail below, in a sufficient amount to achieve target serum concentration. That correlation is also discussed in more detail below.

5 In another embodiment, menopause is treated with precursor as set forth above, in combination with periodic administration of a progestin such as medroxyprogesterone acetate (e.g. Provera) which is preferably administered intermittently, e.g. at a dosage of 2-10 mg per day for 12 consecutive days, said 12-day periods being spaced 20 days to 5 months apart. A combination therapy using a precursor, an estrogen and a progestin may also be used, 10 preferably at the dosages discussed herein for each component.

The sex steroid precursor used in the invention may be administered with or without additional carrier or diluent by the oral route but requires an additional carrier or diluent when administered by the preferred percutaneous 15 or transmucosal route. In a pharmaceutical composition for oral administration, DHEA or other precursor is preferably present in a concentration between 5 and 98% by weight relative to total weight of the composition more preferably between 50 and 98 percent, especially between 80 and 98 percent. If estrogen such as estradiol is present, its concentration 20 is preferably from 0.04 to 0.4 percent by weight. A single precursor such as DHEA may be the only active ingredient, or alternatively, a plurality of precursors and/or their analogues may be used (e.g., a combination of DHEA and DHEA-S, or a combination of two or more compounds converted *in vivo* to DHEA or DHEA-S, or a combination of DHEA and one or more analogues 25 thereof which are converted to DHEA *in vivo*, etc. Where a combination is used, the total dosage of the sum of all precursors should be equal to the dosage range recited above for DHEA used alone, with appropriate adjustment for the different molecular weights of DHEA analogues such as DHEA esters and their conversion rate to DHEA. Thus, if a DHEA ester is 30 used instead of DHEA, dosage should be increased by a multiple equivalent to the ratio of the molecular weight of the DHEA ester to the molecular weight of DHEA. The blood level of DHEA is the final criteria of adequate dosage which takes into account individual variation in absorption and metabolism.

35 Preferably, the attending clinician will, especially at the beginning of treatment, monitor an individual patient's overall response and serum levels

SUBSTITUTE SHEET

- 31 -

of both estrogen and DHEA (in comparison to the preferred serum concentrations discussed above), and monitor the patient's overall response to treatment, adjusting dosages as necessary where a given patients' metabolism or reaction to treatment is atypical. One approach would be to
5 start treating with DHEA or its analogue(s) alone and to add the estrogen only if estrogen blood levels remain too low. Many patients may be treated only with the precursors of the invention without additional estrogen.

Treatment in accordance with the invention is suitable for indefinite
10 continuation. Except for the higher dosage indications discussed above, it is expected that DHEA treatment will simply maintain DHEA levels within a range similar to that which occurs naturally in women before menopause (serum concentration between 4 and 10 micrograms per liter), or naturally in
15 young adult men (serum concentration between 4 and 10 micrograms per liter). Accordingly, undesirable side effects from sustained DHEA treatment are expected to be either minimal or nonexistent. Avoiding side effects from sustained estrogen use may be achieved in ways already known to the art, for example, by intermittent (or in some embodiments continuous)
20 administration of a progestin (e.g., medroxy-progesterone acetate) at a daily oral dose of 2 to 10 mg.

In order to facilitate the combination therapy aspect of the invention, for any indication discussed herein, the invention contemplates pharmaceutical compositions which include both the estrogen and the second active
25 compound (the precursor) in a single composition for simultaneous administration. The composition may be suitable for administration in any traditional manner including but not limited to oral administration, subcutaneous injection or intramuscular injection. In other embodiments, a kit is provided wherein the kit includes the estrogen and second compound
30 (precursor(s)) in separate containers. In addition to other modes of administration, the second compound as well as the estrogen may also be administered transdermally in accordance with the invention as discussed in more detail below. Thus, the kit may include appropriate materials for transdermal administration, e.g., ointments, lotions, gels, creams, sustained
35 release patches and the like. The same strategy applies to the progestin. As

SUBSTITUTE SHEET

- 32 -

noted above, estrogens are disfavored in combination therapies herein involving male patients.

Applicants have discovered that administration of DHEA has utility in the treatment and/or prevention of vaginal atrophy, osteoporosis, skin atrophy, uterine cancer, vaginal cancer, urinary incontinence, hypogonadism and diminished libido, and improves the overall balance of circulating sex steroids, including estrogens and androgens. It is believed that the prior art has not previously suggested that these conditions respond to DHEA treatment. It is believed that DHEA, DHEA-S or a compound converted *in vivo* to either may be useful in the treatment of each of these disorders.

Prior art methods for systemic administration of DHEA have included oral and injection. Because DHEA treatments are often of prolonged and indefinite duration, repeated delivery by injection is very inconvenient. Oral administration, however, has proven relatively inefficient because orally administered DHEA goes first to the liver where a large percentage of it is prevented from entering the general circulation by local degradation.

We have recently observed that DHEA is very efficiently absorbed systemically in both males and females following application to the skin or mucosa (e.g. buccal, vaginal or rectal mucosa). We have discovered that therapeutically efficient doses of DHEA may be administered by the percutaneous or transmucosal route, thus avoiding first passage of the steroid through the liver as results from oral administration, and further avoiding the discomfort and inconvenience of administering DHEA by injection.

Accordingly, the present invention provides delivery systems for the administration of DHEA, DHEA-S and analogous compounds converted *in vivo* to either through the skin or mucosa. These systems are believed to be more efficient than oral administration because the liver is bypassed. These systems are also significantly less painful and more convenient than injections.

35

SUBSTITUTE SHEET

- 33 -

When DHEA, DHEA-S, or analogous compounds converted to DHEA or DHEA-S *in vivo* are formulated for transdermal penetration, any of a number of art-recognized transdermal penetration systems may be utilized. For example, DHEA may be prepared as part of an ointment, lotion, gel or cream for rubbing onto a patient's skin. Active ingredient is preferably present at from 7% to 20% by weight relative to the total weight of the pharmaceutical composition more preferably between 8 and 12%. Alternatively, the active ingredient may be placed into a transdermal patch having structures known in the art, for example, structures such as those set forth in E.P. Patent No.0279982.

When formulated as an ointment, lotion, gel or cream or the like, the active compound is admixed with a suitable carrier which is compatible with human skin or mucosa and which enhances transdermal penetration of the compound through the skin or mucosa. Suitable carriers are known in the art and include but are not limited to Klucel HF and Glaxal base. Some are commercially available, e.g., Glaxal base available from Glaxal Canada Limited Company. Other suitable vehicles can be found in Koller and Buri, S.T.P. Pharma 3(2), 115-124, 1987. The carrier is preferably one in which the active ingredient(s) is (are) soluble at ambient temperature at the concentration of active ingredient that is used. The carrier should have sufficient viscosity to maintain the precursor on a localized area of skin or mucosa to which the composition has been applied, without running or evaporating for a time period sufficient to permit substantial penetration of the precursor through the localized area of skin or mucosa and into the bloodstream where it will cause a measurable and desired increase in serum DHEA concentration. The carrier is typically a mixture of several components, e.g. pharmaceutically acceptable solvents and a thickening agent. A mixture of organic and inorganic solvents can aid hydrophylic and lipophylic solubility, e.g. water and an alcohol such as ethanol.

Desirably, the carrier is one which, if formulated as 10% DHEA and 90% carrier (by weight) and applied twice daily in an amount providing 100 mg of DHEA to the abdominal area, will elevate serum concentration of DHEA in a typical patient by at least 0.35 micrograms per liter per 50 kg of body weight. As mentioned earlier, a Glaxal base, when used as carrier under these

SUBSTITUTE SHEET

- 34 -

conditions, provides an elevation of serum DHEA concentration of about 0.7 µg per liter per 50 kg body weight.

5 The carrier may also include various additives commonly used in ointments and lotions and well known in the cosmetic and medical arts. For example, fragrances, antioxidants, perfumes, gelling agents, thickening agents such as carboxymethylcellulose, surfactants, stabilizers, emollients, coloring agents and other similar agents may be present. When used to treat systemic diseases, the site of application on the skin should be changed in order to
10 avoid excess local concentration of steroids and possible overstimulation of the skin and sebaceous glands by androgenic metabolites of DHEA.

The precursor can also be administered, in some instances, by the oral route, and may be formulated with conventional pharmaceutical excipients, e.g.
15 spray dried lactose and magnesium stearate into tablets or capsules for oral administration at concentrations providing easy dosage in a range from 0.25 to 2.5 grams per day per 50 kg of body weight.

The active substance can be worked into tablets or dragee cores by being
20 mixed with solid, pulverulent carrier substances, such as sodium citrate, calcium carbonate or dicalcium phosphate, and binders such as polyvinyl pyrrolidone, gelatin or cellulose derivatives, possibly by adding also lubricants such as magnesium stearate, sodium lauryl sulfate, "Carbowax" or polyethylene glycol. Of course, taste-improving substances can be added in
25 the case of oral administration forms.

As further forms, one can use plug capsules, e.g. of hard gelatin, as well as closed self-gelatin capsules comprising a softener or plasticizer, e.g. glycerine. The plug capsules contain the active substance preferably in the form of
30 granulate, e.g. in mixture with fillers, such as lactose, saccharose, mannitol, starches, such as potato starch or amylopectin, cellulose derivatives or highly dispersed silicic acids. In self-gelatin capsules, the active substance is preferably dissolved or suspended in suitable liquids, such as vegetable oils or liquid polyethylene glycols.

35

SUBSTITUTE SHEET

- 35 -

The concentration of active ingredient in the ointment, cream, gel or lotion is typically from about 7 to 20 percent preferably between 8 and 12 percent and preferably 10 percent (by weight relative to the total weight of the lotion, cream, gel or ointment). Within the preferred ranges, higher concentrations
5 allow a suitable dosage to be achieved while applying the lotion, ointment, gel or cream to a lesser surface area of the skin than would be possible at lower concentrations and allows more freedom in choosing the body parts to which the ointment or lotion will be applied. For example, it is well known in the art that a compound which is capable of transdermal penetration normally
10 penetrates more efficiently at some points in the body than in others. For example, penetration is very efficient on the forearm and considerably less efficient on the palms.

The lotion, ointment, gel or cream should be thoroughly rubbed into the skin
15 so that no excess is plainly visible, and the skin should not be washed in that region until most of the transdermal penetration has occurred preferably at least 15 minutes and, more preferably, at least 30 minutes.

A transdermal patch may be used to deliver precursor in accordance with
20 known techniques. It is typically applied for a much longer period, e.g., 1 to 4 days, but typically contacts active ingredient to a smaller surface area, allowing a slow and constant delivery of active ingredient.

A number of transdermal drug delivery systems that have been developed,
25 and are in use, are suitable for delivering the active ingredient of the present invention. The rate of release is typically controlled by a matrix diffusion, or by passage of the active ingredient through a controlling membrane.

Mechanical aspects of transdermal devices are well known in the art, and are
30 explained, for example, in United States Patents 5,162,037, 5,154,922, 5,135,480, 4,666,441, 4,624,665, 3,742,951, 3,797,444, 4,568,343, 5,064,654, 5,071,644, 5,071,657, the disclosures of which are incorporated herein by reference. Additional background is provided by European Patent 0279982 and British Patent Application 2185187.

35

SUBSTITUTE SHEET

- 36 -

The device may be any of the general types known in the art including adhesive matrix and reservoir-type transdermal delivery devices. The device may include drug-containing matrixes incorporating fibers which absorb the active ingredient and/or carrier. In a reservoir-type device, the reservoir may
5 be defined by a polymer membrane impermeable to the carrier and to the active ingredient.

In a transdermal device, the device itself maintains active ingredient in contact with the desired localized skin surface. In such a device, the
10 viscosity of the carrier for active ingredient is of less concern than with a cream or gel. A solvent system for a transdermal device may include, for example, oleic acid, linear alcohol lactate and dipropylene glycol, or other solvent systems known in the art. The active ingredient may be dissolved or suspended in the carrier.

15 For attachment to the skin, a transdermal patch may be mounted on a surgical adhesive tape having a hole punched in the middle. The adhesive is preferably covered by a release liner to protect it prior to use. Typical material suitable for release includes polyethylene and polyethylene-coated
20 paper, and preferably silicone-coated for ease of removal. For applying the device, the release liner is simply peeled away and the adhesive attached to the patient's skin. In United States Patent 5,135,480, the disclosure of which is incorporated by reference, Bannon et al. describe an alternative device having a non-adhesive means for securing the device to the skin.

25 Except for the higher dosage indications noted above (e.g. contraception), the target serum concentration of DHEA is comparable regardless of whether sex steroid precursor is being used as part of a combination therapy for treatment of menopause or is being used (by itself or in combination with
30 estrogen and/or progestin) for the treatment of skin deterioration, vaginal atrophy, urinary incontinence, uterine cancer, ovarian cancer, osteoporosis, hypogonadism or diminished libido in accordance with the invention or for the treatment of a wide variety of conditions related to decreased secretion of DHEA by the adrenals. It is pointed out that dosage of DHEA, DHEA-S or
35 any analog discussed herein can all be correlated to a target serum

SUBSTITUTE SHEET

- 37 -

concentration of DHEA because all are converted *in vivo*, either directly or indirectly, into DHEA.

- 5 The percutaneous or transmucosal delivery system of the invention may also be used as a novel and improved delivery system for the prevention and/or treatment of osteoporosis or other diseases which respond favorably to treatment with DHEA. The desired target serum levels for these latter purposes is also the same as indicated above.
- 10 DHEA used for percutaneous or transmucosal application can be in the form of the free alcohol or of one or more of its derivatives, e.g. valerate, benzoate, acetate, enanthate and fatty ester derivatives. The delivery of DHEA or of its analogues through the skin is an acceptable, comfortable and noninvasive way of administering such compound. It also avoids
- 15 gastrointestinal irritation and degradation of the compound and toxicologic problems due to first passage through the liver before reaching the general circulation.
- 20 One method for preventing or inhibiting growth of breast and endometrial carcinoma cells is activation of the androgen receptor with an effective compound having an affinity for the receptor site such that it binds to the androgen receptor at low concentrations while not significantly activating other classes of steroid receptors linked to potential side effects.
- 25 Since DHEA is a natural source of androgens (Labrie, Mol. Cell. Endocrinol. 78: C113-C118, 1991) and the secretion of this compound markedly decreases during aging, its replacement should have minimal unwanted side effects.
- 30 Percutaneous or transmucosal delivery of DHEA in accordance with the invention thus provides a novel method for prevention and therapy of diseases responsive to activation of the androgen receptor, e.g. bone loss, obesity, breast cancer, endometrial cancer, ovarian cancer, urinary incontinence, hypogonadism, loss of libido, loss of muscle mass, loss of
- 35 energy, and other aging processes. The invention is also useful for many diseases wherein activation of the estrogen receptor will have beneficial

SUBSTITUTE SHEET

- 38 -

effects, especially osteoporosis and vaginal atrophy. The invention also represents an improved delivery method (e.g. transdermal) for diseases already in the art as being candidates for DHEA treatment.

5

EXAMPLES OF SOME PREFERRED DERIVATIVES


Derivatives of DHEA or DHEA-S that are expected to convert *in vivo* to DHEA or DHEA-S in accordance with the present invention may be made by the following approaches:

10

- Esterification of the 3 β function into an ester which can be cleaved by esterase (the cleavage does not generate toxic substances) and transformation of the 17-keto group into oxazolidine or thiazolidines which are unstable in the body and regenerate a natural precursor.
- 15 - Formation at position 3 β of an α -acyloxyalkyl ether which can be cleaved by esterase into an unstable hemiacetal. The decomposition of this hemiacetal regenerates the natural precursor.
- Without modification of the 3 β function, transformation of the 17-keto group into oxazolidine or thiazolidines which are unstable in the body and regenerate DHEA or DHEA-S.

20

The compounds disclosed below are expected to convert *in vivo* into dehydroepiandrosterone (DHEA) or dehydroepiandrosterone-sulfate DHEA-S.



25

wherein R³ is hydroxy or sulfate.

SUBSTITUTE SHEET

- 39 -

wherein R^e is selected from the group consisting of hydrogen, benzyl, aryl, straight- or branched-alkyl, straight- or branched-alkenyl and straight- or branched-alkynyl.


5 wherein Z is oxygen or sulfur.

Some derivatives of DHEA being more lipophilic than DHEA itself can be stocked in skin fat and advantageously release DHEA slowly over time.


10 In some preferred compounds of the invention, the function at position 3 is an ester of sulfuric acid (or salts thereof), formic acid, acetic acid, benzoic acid, butyric acid, decanoic acid, enanthic acid, furoic acid, heptanoic acid, isocaproic acid, undecanoic acid, undecylenic acid, palmitic acid, phenylpropionic acid, pivalic acid, propionic acid, valeric acid, carbonic acid (preferably ethylcarbonate or benzylcarbonate).

15

In some preferred compounds of the invention, DHEA (or DHEA-S) is modified by a 17-substituent of the following formula:



20 or




wherein R''' is selected from the group consisting of hydrogen, benzyl, straight- or branched-alkyl and straight- or branched-alkenyl.

25

SUBSTITUTE SHEET

- 40 -

Certain preferred 3 β -ester derivatives of DHEA are listed below:



SUBSTITUTE SHEET


- 41 -

COMPOUNDS	X
dehydroepiandrosterone-3 β -formate	HCO
dehydroepiandrosterone-3 β -acetate	CH ₃ CO
dehydroepiandrosterone-3 β -propionate	CH ₃ CH ₂ CO
dehydroepiandrosterone-3 β -butyrate	CH ₃ (CH ₂) ₂ CO
dehydroepiandrosterone-3 β -valerate	CH ₃ (CH ₂) ₃ CO
dehydroepiandrosterone-3 β -pivalate	(CH ₃) ₃ CCO
dehydroepiandrosterone-3 β -benzoate	C ₆ H ₅ CO
dehydroepiandrosterone-3 β -furoate	C ₄ H ₃ OCO
dehydroepiandrosterone-3 β -cypionate	C ₅ H ₉ (CH ₂) ₂ CO
dehydroepiandrosterone-3 β -lactate	CH ₃ CHOHCO
dehydroepiandrosterone-3 β -decanoate	CH ₃ (CH ₂) ₈ CO
dehydroepiandrosterone-3 β -undecanoate	CH ₃ (CH ₂) ₁₀ CO
dehydroepiandrosterone-3 β -palmitate	CH ₃ (CH ₂) ₁₄ CO
dehydroepiandrosterone-3 β -ethylcarbonate	C ₂ H ₅ OCO
dehydroepiandrosterone-3 β -benzylcarbonate	C ₆ H ₅ CH ₂ OCO
dehydroepiandrosterone-3 β -isocaproate	(CH ₃) ₂ (CH ₂) ₃ CO
dehydroepiandrosterone-3 β -undecylenate	H ₂ C=CH(CH ₂) ₈ CO
dehydroepiandrosterone-3 β -enantate	CH ₃ (CH ₂) ₅ CO
dehydroepiandrosterone-3 β -phenylpropionate	C ₆ H ₅ (CH ₂) ₂ CO
3 β -hydroxymethoxy-5-androsten-17-one acetate	CH ₃ CO ₂ CH ₂
3 β -hydroxymethoxy-5-androsten-17-one decanoate	CH ₃ (CH ₂) ₈ CO ₂ CH ₂

SUBSTITUTE SHEET

- 42 -

Some other preferred DHEA derivatives are listed below:



COMPOUNDS	R'''	Z
3 β -hydroxy-5-androstene-17-spiro(1',3'-thiazolidine-4'-ethyl carboxylate)	C ₂ H ₅	S
3 β -hydroxy-5-androstene-17-spiro(1',3'-thiazolidine-4'-benzyl carboxylate)	C ₆ H ₅ CH ₂	S
3 β -hydroxy-5-androstene-17-spiro(1',3'-thiazolidine-4'-hexyl carboxylate)	C ₆ H ₁₃	S
3 β -hydroxy-5-androstene-17-spiro(1',3'-oxazolidine-4'-ethyl carboxylate)	C ₂ H ₅	O
3 β -hydroxy-5-androstene-17-spiro(1',3'-oxazolidine-4'-benzyl carboxylate)	C ₆ H ₅ CH ₂	O
3 β -hydroxy-5-androstene-17-spiro(1',3'-oxazolidine-4'-hexyl carboxylate)	C ₆ H ₁₃	O

Corresponding 17-substituted analogs of DHEA-S may also be used.

SUBSTITUTE SHEET

- 43 -

EXAMPLES OF SYNTHESIS

5

Example 1

3 β -formyloxy-5-androstene-17-one

Following the procedure described by Ringold (H.J. Ringold, et al., J. Am. Chem. Soc. 78, 816, 1956), dehydroepiandrosterone (2.88 g, 10 mmol) dissolved in 85% formic acid (100 mL) is heated at 60 °C for 1 h. After cooling, the mixture is poured into iced water and after 16 h, crystals are filtered and dried *in vacuo*.

15

Example 2

3 β -acetoxy-5-androstene-17-one

Dehydroepiandrosterone (2.88 g, 10 mmol) is dissolved in a mixture (100 ml) of anhydride acetic and pyridine (1:1 v/v) and left at room temperature for 16 h. The mixture is then poured carefully into iced water and after 16 h, crystals are filtered and dried *in vacuo*.

25

Example 3

Dehydroepiandrosterone-3 β -undecanoate

A solution of undecanoyl chloride (10.2 g, 50 mmol) in CH₂Cl₂ (50 ml) was added to a mixture of 5-androsten-3 β -ol-17-one (11.53 g, 40 mmol), Et₃N (14 ml, 100 mmol) and dimethylaminopyridine (0.6 g, 5 mmol) in CH₂Cl₂ (150 ml) at 5°C. After addition, the mixture was stirred overnight at room temperature. The CH₂Cl₂ layer was washed successively with water, 2N HCl (twice), 5% K₂CO₃ (twice), brine and dried. Removal of the solvent gave the crude product which was recrystallized with a mixture of n-hexane:benzene to give the pure product (13.66 g; 75%), M.P. 84-85°C; ¹H-NMR (CDCl₃); δ 0.83-0.86

SUBSTITUTE SHEET

- 44 -

(m, 6H, C₁₈-CH₃ and CH₃); 1.03 (s, 3H, C₁₉-CH₃); 4.58-4.61 (m, 1H, C₃-H); 5.37 (d, 1H, vinyl, J=4.89 Hz). ¹³C-NMR (CDCl₃) δ : 220.90, 179.25, 139.99, 121.78, 73.38, 51.69, 50.14, 47.49, 38.12, 36.94, 36.72, 35.81, 34.68, 31.87, 31.47, 31.42, 30.76, 29.53, 29.44, 29.27, 29.23, 29.09, 27.73, 25.03, 22.66, 21.86, 20.31, 19.33, 14.09, 13.53.

Example 4

10

3β-acyloxy-5-androstene-17-one

The above esters at position 3β of dehydroepiandrosterone are prepared as follows:

Dehydroepiandrosterone (10 mmol) is dissolved in pyridine (50 mL) and added to a solution of acyl chloride (prepared from the corresponding acid and oxalyl chloride) in the same solvent (50 mL). Dimethylaminopyridine (10%) is then added and the mixture is left at room temperature for 16 h. The mixture is then carefully poured into iced water and extracted with ethyl acetate. The organic phase is washed with diluted HCl, water, saturated sodium bicarbonate and water, dried and evaporated to dryness to give the ester.

25

Example 5

3β-benzyloxycarbonyloxy-5-androsten-17-one

To a stirred solution of dehydroepiandrosterone (2.88 g, 10 mmol) in methylene chloride (100 mL) is added dropwise benzylchloroformate, over a period of 30 min following the known procedure (F. Reber and T. Reichstein, Helv. Chim. Acta, 28, 1164, 1945). After stirring for 3 h, the mixture is washed with water and evaporated to dryness. The residue is then dissolved in acetone and precipitated in iced water. After 16 h, crystals are filtered and dried *in vacuo*.

35

SUBSTITUTE SHEET

- 45 -

Example 6**3 β -ethoxyoxycarbonyloxy-5-androsten-17-one**

- 5 Same procedure as described in example 5 except that ethylchloroformate is used instead of benzylchloroformate.

10

Example 7**3 β -hydroxy-5-androstene-17-spiro-2'-(1', 3'-thiazolidine-4'-ethyl carboxylate)**

- Following the procedure described by Djerassi (C. Djerassi, N. Crossley and
15 M. A. Kielczewski, J. Org. Chem. 27, 1112, 1962), dehydroepiandrosterone (2.88 g, 10 mmol) is dissolved in anhydrous ethanol, sodium acetate is added followed by L-cysteine ethyl ester hydrochloride (18g, 100 mmol) and the mixture is heated overnight under an argon atmosphere. The reaction mixture is then evaporated under vacuum. Methylene chloride is added to
20 precipitate excess of L-cysteine ethyl ester hydrochloride. The solution is then filtered and the filtrate is washed twice with water, dried on magnesium sulfate, filtered and evaporated under vacuum. The residue is triturated with ethanol to give crystals.

25

Example 8**3 β -hydroxy-5-androstene-17-spiro-2'-(1', 3'-thiazolidine-4'-benzyl carboxylate)**

30

Same procedure as described in example 7 except that L-cysteine benzyl ester hydrochloride is used instead of L-cysteine ethyl ester hydrochloride.

35

SUBSTITUTE SHEET.

- 46 -

Example 9

**3 β -hydroxy-5-androstene-17-spiro-2'-(1', 3'-thiazolidine-4'-alkyl
carboxylate)**

- 5 Same procedure as described in example 7 except that different L-cysteine alkyl esters hydrochloride (e.g. L-cysteine hexyl ester hydrochloride) are used instead of L-cysteine ethyl ester hydrochloride.

10

Example 10

**3 β -hydroxy-5-androstene-17-spiro-2'-(1', 3'-oxazolidine-4'-ethyl
carboxylate)**

- 15 The same procedure as described in the Example 7 is used except the oxazolidine derivative is prepared using serine ethyl ester hydrochloride instead of L-cysteine ethyl ester hydrochloride.

20

Example 11

**3 β -hydroxy-5-androstene-17-spiro-2'-(1', 3'-oxazolidine-4'-benzyl
carboxylate)**

- 25 Same procedure as described in example 10 except that serine benzyl ester hydrochloride is used instead of serine ethyl ester hydrochloride.

30

Example 12

**3 β -hydroxy-5-androstene-17-spiro-2'-(1', 3'-oxazolidine-4'-alkyl
carboxylate)**

- 35 Same procedure as described in example 10 except that different serine alkyl esters hydrochloride (e.g. serine hexyl ester hydrochloride) are used instead of serine ethyl ester hydrochloride.

SUBSTITUTE SHEET

Example 13

5

3 β -hydroxymethoxy-5-androsten-17-one acetate

To a solution of dehydroepiandrosterone (2.88 g, 10 mmol) in THF (100 mL) is added sodium hydride (11 mmol, 60% in oil) at room temperature under an argon atmosphere. When all the sodium hydride has reacted, chloromethyl acetate (prepared from acetyl chloride and formaldehyde (or derivative) using ZnCl₂ as catalyst) is added and the mixture is heated for a few hours. After cooling, the mixture is poured into water and extracted with ethyl acetate. The organic phase is then washed with water, dried, filtered and evaporated to dryness to give the desired compound.

15

EXAMPLES OF PHARMACEUTICAL COMPOSITIONS

In one aspect, the present invention involves applying DHEA (or its analogues) formulation to the skin or other epithelial tissue for a time period sufficient to permit sufficient penetration of the compound for systemic or topical action, as desired. The composition may be applied as a gel, a cream, an ointment, a lotion or the like or may involve use of a delivery system as described in U.S. Patent Nos 3,742,951, 3,797,494 or 4,568,343. Devices as described in U.S. Patent Nos 5,064,654, 5,071,644 or 5,071,657 can also be used to facilitate steroid absorption.

All the pharmaceutical compositions of the present invention may contain appropriate preservatives known in the art.

30

The following non-limiting examples describe the preparation of a typical cream, lotion, gel and ointment, respectively. In addition to these vehicles, one skilled in the art may choose other vehicles in order to adapt to specific dermatologic needs.

35

- 48 -

Example 14

A typical lotion contains (W/W) 10% DHEA, 15% propylene glycol and 70%
5 ethanol and water 5%.

Example 15

10 A typical gel contains (W/W) 10% DHEA, 5% propylene glycol, 0.2%
Carbomer 940 (available as Carbopol 940^R from B.F. Goodrich), 40% water,
0.2% triethanolamine, 2% PPG-12-Buteh-16 (available as Ucon^R fluid 50
15 from Union Carbide), 1% hydroxypropyl and 41.6% ethanol (95% ethanol-5%
water).

Example 16

20 A typical ointment contains (W/W) 10% DHEA, 13% propylene glycol, 74%
petrolatum, 2.9% glycerylmonostearate and 0.1% polyparaben.

25

Example 17

A typical cream contains (W/W) 10% DHEA, 0.2% propylparaben, 5% lanolin
oil, 7.5% sesame oil, 5% cetyl alcohol, 2% glyceryl monostearate, 1%
30 triethanolamine, 5% propylene glycol, 0.1% Carbomer 940^R and 64.2%
water.

In each of the foregoing Examples 14-17, a progestin and/or an estrogen may
be added. For example 0.005 to 0.02% 17 β -estradiol and/or 0.2 to 2.0%
35 medroxyprogesterone acetate may be added with corresponding reductions
in water or ethanol or petrolatum. DHEA permeability can be enhanced by

SUBSTITUTE SHEET

- 49 -

various techniques in order to reduce the dose used. Methods and compositions for enhancing permeability of an active compound can be found, for example, in U.S. Patents Nos 5,051,260, 4,006,218, 3,551,554, 3,472,931, 4,568,343, 3,989,816 and 4,405,616.

5

Although the present invention has been described in relation to particular embodiments thereof, many other variations and modifications and other uses will become apparent to those skilled in the art. It is preferred, therefore, that the present invention be limited not by the specific disclosure

10 herein, but only by the appended claims.

SUBSTITUTE SHEET.

- 50 -

WHAT IS CLAIMED IS:

1. A method for preventing or treating symptoms of menopause comprising administering to a patient in need of such prevention or treatment, with or without a pharmaceutical carrier or diluent, at least one
5 sex steroid precursor selected from the group consisting of dehydroepiandrosterone, dehydroepiandrosterone-sulfate and compounds converted *in vivo* to either of the foregoing, said method further comprising administering at least one additional agent selected from the group consisting of an estrogen and a progestin, as part of a combination therapy.
10
2. A pharmaceutical composition comprising a sex steroid precursor selected from the group consisting of DHEA, DHEA-S, and compounds converted *in vivo* to either, said composition further comprising a supplemental agent selected from the group consisting of an estrogen and a
15 progestin.
3. The pharmaceutical composition of claim 2 wherein said precursor is DHEA.
- 20 4. The pharmaceutical composition of claim 2 further comprising a pharmaceutically acceptable diluent or carrier.
5. The pharmaceutical composition of claim 2 wherein said composition comprises a pharmaceutically acceptable diluent or carrier for
25 percutaneous or transmucosal administration, wherein said estrogen is present at a concentration between 0.005 and 1.0 percent by weight relative to the total weight of the pharmaceutical composition and wherein said precursor is present at a concentration between 7.0 and 20 percent by weight.
30

SUBSTITUTE SHEET.

- 51 -

6. A kit comprising a first container having therein at least one sex steroid precursor selected from the group consisting of dehydroepiandrosterone, dehydroepiandrosterone sulphate, and compounds converted *in vivo* to either of the foregoing and further comprising at least one
5 additional container having therein at least one active ingredient selected from the group consisting of an estrogen and a progestin.

7. The kit of claim 6 wherein said kit provides each of a progestin, an estrogen and a sex steroid precursor.

10

8. A method of preventing or treating vaginal atrophy comprising administering to a patient in need of such prevention or treatment, with or without pharmaceutical diluent or carrier, an effective amount of at least one sex steroid precursor selected from the group consisting of
15 dehydroepiandrosterone, dehydroepiandrosterone sulphate, and compounds converted *in vivo* to either of the foregoing.

9. A method of preventing or treating hypogonadism comprising administering to a patient in need of such prevention or treatment,
20 with or without pharmaceutical diluent or carrier, an effective amount of at least one sex steroid precursor selected from the group consisting of dehydroepiandrosterone, dehydroepiandrosterone sulphate, and compounds converted *in vivo* to either of the foregoing.

25 10. A method of preventing or treating diminished libido administering to a patient in need of such prevention or treatment, with or without pharmaceutical diluent or carrier, an effective amount of at least one sex steroid precursor selected from the group consisting of dehydroepiandrosterone, dehydroepiandrosterone sulphate, and compounds
30 converted *in vivo* to either of the foregoing.

SUBSTITUTE SHEET.

- 52 -

11. A method of preventing or treating skin atrophy comprising administering to a patient in need of such prevention or treatment, with or without pharmaceutical diluent or carrier, an effective amount of at least one sex steroid precursor selected from the group consisting of dehydroepiandrosterone, dehydroepiandrosterone sulphate, and compounds converted *in vivo* to either of the foregoing.

12. A therapeutic method of treating reduced or imbalanced concentrations of sex steroids comprising applying an effective amount of a pharmaceutical composition for percutaneous or transmucosal delivery to an outer surface of skin or mucosa of a patient in need of such treatment, said pharmaceutical composition comprising a carrier having therein at least one sex steroid precursor selected from the group consisting of dehydroepiandrosterone, dehydroepiandrosterone sulphate, and compounds converted *in vivo* to either of the foregoing, said precursor being present at a concentration of at least 7% by weight relative to the total pharmaceutical composition, and said carrier being compatible with skin or mucosa and allowing penetration of said precursor through said skin or mucosa, said carrier having sufficient viscosity to maintain said precursor on a localized area of skin or mucosa, without running or evaporating, for a time period sufficient to permit substantial penetration of said precursor through said localized area of said skin or mucosa.


13. The method of claim 12 where the method is used for prevention or treatment of an indication selected from the group consisting of obesity, cardiovascular disease, atherosclerosis, breast cancer, endometrial cancer, loss of muscle mass, diabetes, fatigue or loss of energy, connective tissue diseases, loss of memory, and symptoms of menopause.

SUBSTITUTE SHEET

- 53 -

14. A pharmaceutical composition for percutaneous or transmucosal delivery, said pharmaceutical composition comprising a carrier having therein at least one sex steroid precursor selected from the group consisting of dehydroepiandrosterone, dehydroepiandrosterone sulphate, and compounds converted *in vivo* to either of the foregoing, said precursor being present at a concentration of at least 7% by weight relative to the total pharmaceutical composition, and said carrier being compatible with skin or mucosa and allowing penetration of said precursor through said skin or mucosa, said carrier having sufficient viscosity to maintain said precursor on a localized area of skin or mucosa, without running or evaporating, for a time period sufficient to permit substantial penetration of said precursor through said localized area of said skin or mucosa.

15. A compound having the following formula:




- 20 wherein X is selected from the group consisting of H, $\text{R}-\overset{\text{O}}{\parallel}{\text{C}}-$, $\text{RCO}_2\text{CHR}^a-$ and R^bSO_2- (R being selected from the group consisting of hydrogen, straight- or branched-alkyl, straight- or branched-alkenyl, straight- or branched-alkynyl, aryl, furyl, straight- or branched-alkoxy, straight- or branched-alkenyloxy, straight- or branched-alkynyloxy, aryloxy, furyloxy and halogeno analogs of the foregoing; R^a being hydrogen or (C_1-C_6) alkyl; and, R^b being selected from the group consisting of hydroxyl (or salts thereof), methyl, phenyl and p-toluy!);

SUBSTITUTE SHEET

- 54 -

wherein Y is a divalent substituted or unsubstituted moiety of the formula:




5

(Z being an oxygen or sulfur atom), and wherein Y and the carbon atom to which it is bonded together form a closed saturated 5-membered ring.

16. A pharmaceutical composition comprising a
 10 pharmaceutically acceptable diluent or carrier and an amount of at least one compound of claim 15 effective to raise serum concentration of DHEA or DHEA-S in a patient to whom said composition is administered.

17. A compound having the formula:

15



- wherein R' is selected from the group consisting of C₃-C₂₀ straight- or branched-alkyl, C₃-C₂₀ straight- or branched-alkenyl, C₃-C₂₀ straight- or
 20 branched-alkynyl, aryl and halo-substituted analogs of the foregoing.

18. A pharmaceutical composition comprising
 pharmaceutically acceptable diluent or carrier and an amount of at least one


SUBSTITUTE SHEET

- 55 -

compound of claim 17 effective to raise serum concentration of DHEA or DHEA-S in a patient to whom said composition is administered.

19. A compound of the formula:

5



R^d being selected from the group consisting of hydrogen, straight- or branched-alkyl, straight- or branched-alkenyl, straight- or branched-alkynyl, aryl, furyl, straight- or branched-alkoxy, straight- or branched-alkenyloxy, straight- or branched-alkynyloxy, aryloxy, furyloxy and halogeno analogs of the foregoing.

wherein R_a is a hydrogen or (C_1-C_6) alkyl.

15


20. A pharmaceutical composition comprising a pharmaceutically acceptable diluent or carrier and an amount of at least one compound of claim 19 effective to raise serum concentration of DHEA or DHEA-S in a patient to whom said composition is administered.

20

21. A compound having the formula:

SUBSTITUTE SHEET

- 56 -



wherein R^3 is hydroxy or sulfate.

- 5 wherein R^e is selected from the group consisting of hydrogen, benzyl, aryl, straight- or branched-alkyl, straight- or branched-alkenyl and straight- or branched-alkynyl.

wherein Z is oxygen or sulfur.

10

22. A pharmaceutical composition comprising a pharmaceutically acceptable diluent or carrier and an amount of at least one compound of claim 21 effective to raise serum concentration of DHEA or DHEA-S in a patient to whom said composition is administered.

15

23. A transdermal patch which continuously delivers an active ingredient selected from the group consisting of dehydroepiandrosterone, dehydroepiandrosterone sulphate and analogues which are converted *in vivo* to either said patch having a means for attaching skin, a receptical for said active ingredient and a means for conducting said active ingredient into contact with skin of a user.
- 20

24. A method for the prevention or treatment of osteoporosis comprising administering to a patient in need of such prevention or treatment with or without a pharmaceutical diluent or carrier, a therapeutically effective
- 25

SUBSTITUTE SHEET

- 57 -

amount of a sex steroid precursor selected from the group consisting of DHEA, DHEA-S, and compounds that are converted *in vivo* to either.

25. A method for the prevention or treatment of urinary
5 incontinence comprising administering to a patient in need of such prevention or treatment, with or without pharmaceutical carrier or diluent, a therapeutically effective amount of a sex steroid precursor selected from the group consisting of DHEA, DHEA-S, and compounds converted *in vivo* to either.

10


26. A method of contraception comprising administering to a female in need of contraception, with or without pharmaceutical carrier or diluent, an effective amount of a sex steroid precursor selected from the group consisting of DHEA, DHEA-S, and compounds converted *in vivo* to
15 either.

27. A method for prevention of ovarian cancer comprising administering to a female patient in need of such prevention, with or without
20 pharmaceutical carrier or diluent, a therapeutically effective amount of an agent selected from the group consisting of DHEA, DHEA-S, and compounds converted *in vivo* to either.

28. A method for prevention or treatment of uterine cancer
25 comprising administering to a female patient in need of such prevention or treatment, with or without pharmaceutical diluent or carrier, a therapeutically effective amount of a sex steroid precursor selected from the group consisting of DHEA, DHEA-S, and compounds converted *in vivo* to either.

SUBSTITUTE SHEET

1/12



** ,p<0.01: Intact versus Int + EM-760


tt, p< 0.01: Ovx versus Ovx + EM-760

FIG. 1

SUBSTITUTE SHEET


2/12

FIGURE 2



3/12

FIGURE 3



SUBSTITUTE SHEET

4/12




FIG. 4

SUBSTITUTE SHEET

5/12




FIG. 5a




FIG. 5b



FIG. 5c




FIG. 5d

SUBSTITUTE SHEET

6/12




FIG. 6a




FIG. 6b




FIG. 6c




FIG. 6d




FIG. 6e




FIG. 6f

SUBSTITUTE SHEET

7/12




FIG.7a




FIG.7b




FIG.7c

SUBSTITUTE SHEET

8/12



FIG.7d




FIG.7e




FIG.7f

SUBSTITUTE SHEET

9/12




FIG. 8a




FIG. 8b




FIG. 8c

SUBSTITUTE SHEET.

10/12



FIG.8d




FIG.8e




FIG.8f

SUBSTITUTE SHEET

11/12




FIG.9a



FIG.9b




FIG.9c

SUBSTITUTE SHEET

12/12




FIG. 9d



FIG. 9e



FIG. 9f

SUBSTITUTE SHEET.

**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ **BLACK BORDERS**
- ☐ **IMAGE CUT OFF AT TOP, BOTTOM OR SIDES**
- ☐ **FADED TEXT OR DRAWING**
- ☐ **BLURRED OR ILLEGIBLE TEXT OR DRAWING**
- ☐ **SKEWED/SLANTED IMAGES**
- ☐ **COLOR OR BLACK AND WHITE PHOTOGRAPHS**
- ☐ **GRAY SCALE DOCUMENTS**
- ☐ **LINES OR MARKS ON ORIGINAL DOCUMENT**
- ☐ **REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY**
- ☐ **OTHER: _____**

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.